

Dorothee PERREAU

Digitally signed  
by Dorothee  
PERREAU

Date:

2026.09.11

11:19:38 +02'00'



3 666560 000019  
G.T.I.N. 036665600000019  
Lot number :  
EXP:

260050/1A  
04/2030

102792BT26-2

Active ingredients per ml:

Activated charcoal 300 mg, Sodium sulfate 62.5 mg

Inactive ingredients:

water, glycerin, potassium sorbate, carboxymethylcellulose sodium, xanthan gum.

Indications:

For use as a digestive adsorbent and cathartic to support and facilitate elimination of undesirable ingested substances following accidental exposure in dogs and cats.




Directions for use:

For oral use only as a **one-time administration** for dogs and cats.

Administer the amount of product according to the following table

- One dot (●) delivers 6 mL (0.02 oz) of product.
- Ten dots (●) correspond to 1 full syringe (60 mL or 2.03 oz).

For animals over 20kg in body weight, use the appropriate combination of syringes. For example, a 30kg animal would receive one full syringe and 5 dots of another syringe.



1. Before first use, wind ring as shown on image ❶ , until the desired number of dots is visible between the barrel and the ring as seen on image ❷.
2. Remove the cap and administer the product directly into the animal's mouth on the back of the tongue (image ❸).
3. After use, clean the syringe tip and replace the cap to prevent the product from drying and blocking the syringe.



Carbodote™  
Prime

ACTIVATED CHARCOAL AND SODIUM SULFATE GEL  
FOR DOGS AND CATS



DP | DÔMES  
PHARMA

Net contents | **60 mL**  
(2.03 oz)

Storage:

Store at room temperature and in a dry place.  
Discard 1 month after opening

Cautions:

CARBODOTE PRIME™ is intended for one time administration. When repeated administration of an adsorbent is necessary, follow up with CARBODOTE REPEAT™. For animal use only.

**Do not administer by mouth to animals unable to swallow.**

**KEEP OUT OF REACH OF CHILDREN.**

Should not be given in conjunction with oral medications as the charcoal may adsorb the therapeutic agent. Allow about 3 hours between the administration of CARBODOTE PRIME and a meal and/or oral treatments. Activated charcoal may cause the stools to turn black and can stain textiles and contact materials.

These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

NDC number: 86189-001-01

Manufactured for:

DOMES PHARMA Inc.

Salt Lake City, UT 84115

1-833-224-2015

info.us@domespharma.com

Carbodote™

Prime

DP | DÔMES  
PHARMA