

Active ingredients per ml:

Activated charcoal 300 mg

Inactive ingredients:

water, glycerin, potassium sorbate, xanthan gum.

Indications:

For use as a digestive adsorbent following accidental ingestion of undesirable substances in dogs and cats.


Directions for use:

For oral use only for dogs and cats.

Administration can be repeated at 4-6 hours intervals for up to 24 hours or as recommended by your veterinarian. Administer the amount of product according to the following table

- One dot (•) delivers to 6 mL (0,02 oz) of product.
 - Ten dots (•) correspond to 1 full syringe (60 mL or 2.03 oz).
- For animals over 20kg in body weight, use the appropriate combination of syringes. For example, a 30kg animal would receive one full syringe and 5 dots of another syringe.

Animal's weight (kg)	Animal's weight (lb)	Number of dots (•)
≤ 2	0.0 - 4.5	1 •
2.1 - 4	4.6 - 8.9	2 ••
4.1 - 6	9.0 - 13.3	3 •••
6.1 - 8	13.4 - 17.8	4 ••••
8.1 - 10	17.9 - 22.2	5 •••••
10.1 - 12	22.3 - 26.6	6 ••••••
12.1 - 14	26.7 - 31.0	7 •••••••
14.1 - 16	31.1 - 35.4	8 ••••••••
16.1 - 18	35.5 - 39.8	9 •••••••••
18.1 - 20	39.9 - 44.1	10 ••••••••••



1. Before first use, wind ring as shown on image ❶, until the desired number of dots is visible between the barrel and the ring as seen on image ❷.
2. Remove the cap. Slightly raise the animal's head. Insert the syringe tip between the cheek and the teeth, towards the back of the mouth to make swallowing easier. Depress the plunger slowly to give the animal time to swallow. Keep the mouth closed for a few seconds if necessary to help the animal swallow (image ❸).
3. After use, clean the syringe tip and replace the cap to prevent the product from drying and blocking the syringe.



Carbodote™ Repeat



ACTIVATED CHARCOAL GEL FOR DOGS AND CATS

3 666560 027931
G.T.I.N. 03666560027931
Lot number :
EXP:
250904/1A
03/2030

102793BT26-2

Net contents (2.03 oz) | 60 mL



Storage:

Store at room temperature and in a dry place. Discard 1 month after opening.

Cautions:

In animals where a cathartic effect is indicated, administer an initial dose of CARBODOTE PRIME™ to facilitate the elimination of toxins.

For animal use only.

Do not administer by mouth to animals unable to swallow. In all cases of suspected poisonings, consult your veterinarian for complete diagnosis and specific treatment.

KEEP OUT OF REACH OF CHILDREN.

Should not be given in conjunction with oral medications as the charcoal may adsorb the therapeutic agent. Allow about 3 hours between the administration of CARBODOTE REPEAT™ and a meal and/or oral treatments. Activated charcoal may cause the stools to turn black and can stain textiles and contact materials.

These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

NDC number: 86189-002-01

Manufactured for:

DOMES PHARMA Inc.
Salt Lake City, UT 84115
1-833-224-2015
info.us@domespharma.com

Carbodote® Repeat

DP DOMES PHARMA