

CARBODOTE REPEAT- activated charcoal gel

Domes Pharma

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click [here](#).

CARBODOTE REPEAT

Active Ingredients per ml:

Activated charcoal 300 mg

Inactive ingredients:

water, glycerin, potassium sorbate, xanthan gum

Indications:


For use as a digestive adsorbent following accidental ingestion of undesirable substances in dogs and cats.

Directions for use:

For oral use only for dogs and cats.

Administration can be repeated at 4-6 hour intervals for up to 24 hours or as recommended by your veterinarian. Administer the amount of product according to the following table. One dot delivers 6mL (0.02oz) of product. Ten dots corresponds to 1 full syringe (60mL or 2.03oz). For animals over 20kg in body weight, use the appropriate combination of syringes. For example, a 30kg animal would receive one full syringe and 5 dots of another syringe.

Animal's weight (kg)	Animal's weight (lb)	Number of dots (•)
≤ 2	0.0 - 4.5	1 •
2.1 - 4	4.6 - 8.9	2 ••
4.1 - 6	9.0 - 13.3	3 •••
6.1 - 8	13.4 - 17.8	4 ••••
8.1 - 10	17.9 - 22.2	5 •••••
10.1 - 12	22.3 - 26.6	6 ••••••
12.1 - 14	26.7 - 31.0	7 •••••••
14.1 - 16	31.1 - 35.4	8 ••••••••
16.1 - 18	35.5 - 39.8	9 •••••••••
18.1 - 20	39.9 - 44.1	10 ••••••••••



1. Before first use, wind ring as shown on image ❶, until the desired number of dots is visible between the barrel and the ring as seen on image ❷.
2. Remove the cap. Slightly raise the animal's head. Insert the syringe tip between the cheek and the teeth, towards the back of the mouth to make swallowing easier. Depress the plunger slowly to give the animal time to swallow. Keep the mouth closed for a few seconds if necessary to help the animal swallow (image ❸).
3. After use, clean the syringe tip and replace the cap to prevent the product from drying and blocking the syringe.

Storage:

Store at room temperature and in a dry place.

Discard 1 month after opening.

Cautions:

In animals where a cathartic effect is indicated, administer an initial dose of CARBODOTE PRIME to facilitate the elimination of toxins.

For use in animals only.

Do not administer by mouth to animals unable to swallow.

In all cases of suspected poisonings, consult your veterinarian for complete diagnosis and specific treatment.

KEEP OUT OF REACH OF CHILDREN.

Should not be given in conjunction with oral medications as the charcoal may adsorb the therapeutic agent. Allow about 3 hours between the administration of CARBODOTE REPEAT and a meal and/or oral treatments. Activated charcoal may cause the stools to turn black and can stain textiles and contact materials.

These statements have not been evaluated by the Food and Drug Administration. This

product is not intended to diagnose, treat, cure or prevent any disease.

CARBODOTE REPEAT Label

Syringe Label

102793EA26-1 - 1129/02M

Active ingredients per mL: Activated charcoal 300 mg.
Inactive ingredients: Water, glycerin, potassium sorbate, xanthan gum.

Indications: For use as a digestive adsorbent following accidental ingestion of undesirable substances in dogs and cats.

Directions for use: For oral use only.
See carton box for complete instructions and cautions.

Cautions: For animal use only. Keep out of reach of children.

Storage: Store at room temperature and in a dry place.
Discard 1 month after opening.

NDC Number: 86189-002-01

Manufactured for:
DOMES PHARMA Inc.
Salt Lake City, UT 84115
1-833-224-2015
info.us@domespharma.com

Carbodote™
Repeat

ACTIVATED CHARCOAL GEL FOR DOGS AND CATS 

DP | DOMES PHARMA Net contents (2.03 oz) | **60 mL**

Box Label

Active ingredients per mL:
Activated charcoal 300 mg

Inactive ingredients:
water, glycerin, potassium sorbate, xanthan gum.


Indications:
For use as a digestive adsorbent following accidental ingestion of undesirable substances in dogs and cats.

Directions for use:
For oral use only for dogs and cats.

Administration can be repeated at 4-6 hours intervals for up to 24 hours or as recommended by your veterinarian. Administer the amount of product according to the following table

- One dot (•) delivers to 6 mL (0.02 oz) of product.
 - Ten dots (•) correspond to 1 full syringe (60 mL or 2.03 oz).
- For animals over 20kg in body weight, use the appropriate combination of syringes. For example, a 30kg animal would receive one full syringe and 5 dots of another syringe.

Animal's weight (kg)	Animal's weight (lb)	Number of dots (•)
≤ 2	0.0 - 4.5	1 •
2.1 - 4	4.6 - 8.9	2 ••
4.1 - 6	9.0 - 13.3	3 •••
6.1 - 8	13.4 - 17.8	4 ••••
8.1 - 10	17.9 - 22.2	5 •••••
10.1 - 12	22.3 - 26.6	6 ••••••
12.1 - 14	26.7 - 31.0	7 •••••••
14.1 - 16	31.1 - 35.4	8 ••••••••
16.1 - 18	35.5 - 39.8	9 •••••••••
18.1 - 20	39.9 - 44.1	10 ••••••••••



1. Before first use, wind ring as shown on image 1, until the desired number of dots is visible between the barrel and the ring as seen on image 2.
2. Remove the cap. Slightly raise the animal's head. Insert the syringe tip between the cheek and the teeth, towards the back of the mouth to make swallowing easier. Depress the plunger slowly to give the animal time to swallow. Keep the mouth closed for a few seconds if necessary to help the animal swallow (image 3).
3. After use, clean the syringe tip and replace the cap to prevent the product from drying and blocking the syringe.



Carbodote™
Repeat.

ACTIVATED CHARCOAL GEL FOR DOGS AND CATS



DP

Net contents (2.03 oz) **60 mL**

Storage:
Store at room temperature and in a dry place.
Discard 1 month after opening.

Cautions:
In animals where a cathartic effect is indicated, administer an initial dose of CARBODOTE PRIME™ to facilitate the elimination of toxins.

For animal use only.

Do not administer by mouth to animals unable to swallow.

In all cases of suspected poisonings, consult your veterinarian for complete diagnosis and specific treatment.

KEEP OUT OF REACH OF CHILDREN.

Should not be given in conjunction with oral medications as the charcoal may adsorb the therapeutic agent. Allow about 3 hours between the administration of CARBODOTE REPEAT™ and a meal and/or oral treatments. Activated charcoal may cause the stools to turn black and can stain textiles and contact materials.

These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

NDC number: 86189-002-01

Manufactured for:

DOMES PHARMA Inc.

Salt Lake City, UT 84115

1-833-224-2015

info.us@domespharma.com

CARBODOTE REPEAT

activated charcoal gel

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:86189-002
Route of Administration	oral		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACTIVATED CHARCOAL (UNII: 2P3VWU3H10) (ACTIVATED CHARCOAL - UNII:2P3VWU3H10)	ACTIVATED CHARCOAL	300 mg in 1 mL

Inactive Ingredients				
Ingredient Name				Strength
WATER (UNII: 059QF0KO0R)				
GLYCERIN (UNII: PDC6A3C0OX)				
POTASSIUM SORBATE (UNII: 1VPU26JZZ4)				
XANTHAN GUM (UNII: TTV12P4NEE)				
Product Characteristics				
Color	black	Score		
Shape		Size		
Flavor		Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:86189-002-01	60 mL in 1 SYRINGE		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
unapproved drug other			07/06/2026	

Labeler -
Domes Pharma (770769404)

Establishment			
Name	Address	ID/FEI	Business Operations
D2M Sante		569606767	manufacture, pack, label, analysis, api manufacture